

# 4 Renormalization group (RG)

## 4.1 Concept of RG

Assumption: The relevant physics describing phases and phase transition is governed by the behavior at large length scales ( $\hat{=}$  low energy)

$$L \gg a \leftarrow \begin{matrix} \text{microscopic} \\ \text{length scale} \\ \uparrow \\ \text{typical length} \\ \text{scale} \end{matrix}$$

Example (magnet):  $\langle S_i^z S_{i+1}^z \rangle \neq 0$  for all  $T$

$$\lim_{|i-j| \rightarrow \infty} \langle S_i^z S_j^z \rangle \begin{cases} = 0 & \text{for } T > T_c \\ \neq 0 & \text{for } T < T_c \end{cases}$$

Idea: Successively integrate out short-distance ( $\hat{=}$  high-energy) modes to obtain an effective theory at large length scales ( $\hat{=}$  low energy)

$$\text{E.g.: } S(g_1, g_2, \dots) \mapsto S(g'_1, g'_2, \dots)$$


$\uparrow$  action       $\uparrow$  couplings       $\uparrow$  if  $S$  is sufficiently general

$\Rightarrow$  couplings become scale-dependent  $g_i \rightarrow g_i(L)$

RG flow:


Change of couplings under the successive integration of modes

RG flow diagram (example):



Integrate out high-energy modes with momenta:

$$\frac{\Lambda}{b} \leq |\vec{k}| < \Lambda$$



"momentum shell"

Infinitesimal RG step:

$$b = e^{dt} \text{ with } dt \ll 1, \quad t = \int_0^t dt' \text{ "RG time"}$$

RG flow:

$$\frac{dg_i}{dt} = \beta_i(g_i)$$

↑  
beta functions

Fixed point:

$$\left. \frac{dg_i}{dt} \right|_{g_i^*} = 0$$

Linearized RG flow (around Gaussian fixed point  $g^* = 0$  in theory with one coupling  $g$ ):

$$\beta(g) = \theta g + O(g^2) \quad \text{with } \theta \equiv \text{dim}[g] = \text{const.} \quad \text{"scaling dimension of } g\text{"}$$

Integrated flow: ↑  
constant term vanishes  
since  $g=0$  is a fixed point

$$\frac{dg}{dt} = \theta g \quad \Rightarrow \quad g(t) = g(0) e^{\theta t}$$

Classification of couplings:

$$\text{dim}[g] > 0$$

"relevant coupling"

$g$  increases in RG time

$$\text{dim}[g] < 0$$

"irrelevant coupling"

$g$  decreases in RG time

$$\text{dim}[g] = 0$$

"marginal coupling"

higher-order terms decide its fate  
(marginally relevant, marginally irrelevant,  
or exactly marginal)

# Classification of fixed points:

"stable fixed point": all couplings irrelevant near fixed point

"critical fixed point": exactly one relevant direction

"multicritical fixed point": number of relevant directions  $2 \leq n \leq n_0$   
where  $n_0$  is the number of tuning parameters

"unstable fixed point": number of relevant directions  $n > n_0$

## 4.3 Momentum-shell RG for the $O(N)$ model

Action ( $O(N)$  model):

$$S = \int d^d \vec{x} \left[ \frac{1}{2} (\vec{\nabla} \phi^a(\vec{x}))^2 + \frac{u}{2} (\phi^a(\vec{x}))^2 + \frac{u}{4!} (\phi^a(\vec{x}))^2 \right], \quad a=1, \dots, N$$

↙ tuning parameter ("mass")  
↗ self-interaction coupling

$$= \int_0^\Lambda \frac{d^d \vec{k}}{(2\pi)^d} \frac{1}{2} \phi^a(\vec{k}) (\vec{k}^2 + \dots) \phi^a(\vec{k}) + \frac{u}{4!} \int_0^\Lambda \frac{d^d \vec{k}_1 d^d \vec{k}_2 d^d \vec{k}_3}{(2\pi)^{3d}} \phi^a(\vec{k}_1) \phi^a(\vec{k}_2) \phi^b(\vec{k}_3) \phi^b(-\vec{k}_1 - \vec{k}_2 - \vec{k}_3)$$

where we have rescaled  $\sum_0^2 (\phi^a)^2 \mapsto \phi^a$  and introduced an ultraviolet momentum cutoff  $\Lambda$ ,  $0 \leq |\vec{k}| \leq \Lambda$ , with, e.g.,  $\Lambda \sim \frac{\pi}{a}$  ( $a$ : lattice constant).

Three stages of RG transformation:

1. Eliminate "fast" modes  $\phi_s$  with momenta  $\frac{\Lambda}{b} \leq |\vec{k}| < \Lambda$  ("momentum shell").

$$\phi(\vec{k}) \equiv \underbrace{\Theta\left(\frac{\Lambda}{b} - |\vec{k}|\right)}_{\substack{\uparrow \\ \text{slow modes} \\ 0 \leq |\vec{k}| < \frac{\Lambda}{b}}} \phi_s(\vec{k}) + \underbrace{\Theta\left(|\vec{k}| - \frac{\Lambda}{b}\right)}_{\substack{\uparrow \\ \text{fast modes} \\ \frac{\Lambda}{b} \leq |\vec{k}| < \Lambda}} \phi_s(\vec{k})$$

2. Rescale momenta  $\vec{k} \mapsto \vec{k}' = b\vec{k}$  with  $0 \leq |\vec{k}'| < \Lambda$  for slow modes.

3. Introduce "renormalized" fields  $\phi'(\vec{k}') = b^\gamma \phi_s(\vec{k}'/b)$  with  $\gamma$  chosen such that the new action in terms of  $\phi'$  has the same coefficient for a certain quadratic term.

10.5.24

RG for Gaussian model ( $u=0$ ):

1. Mode elimination:

$$Z = \int \mathcal{D}\phi_s \int \mathcal{D}\phi_s e^{-S_0[\phi_s, \phi_s]}$$

with

$$S_0[\phi_s, \phi_s] = \int_0^{\Lambda/b} \frac{d^d k}{(2\pi)^d} \frac{1}{2} \phi_s(-\vec{k})(\vec{k}^2 + \tau)\phi_s(\vec{k}) + \int_{\Lambda/b}^{\Lambda} \frac{d^d k}{(2\pi)^d} \frac{1}{2} \phi_s(-\vec{k})(\vec{k}^2 + \tau)\phi_s(\vec{k})$$

Thus:

$$Z = \int \mathcal{D}\phi_s e^{-\underbrace{\int_0^{\Lambda/b} \phi_s(k^2 + \tau)\phi_s}_{\equiv S_{\text{eff}}}} \cdot \text{const.}$$

↑ independent of  $\phi_s$

$$\left[ Z_{0s} = \int \mathcal{D}\phi_s e^{-S_{0s}} = \left[ \det(k^2 + \tau) \right]_{\frac{\Lambda}{b} \leq k < \Lambda}^{-1/2} \right]$$

2. Momentum rescaling:

$$S_{\text{eff}} = \int_0^{1/b} \frac{d^d \vec{k}}{(2\pi)^d} \frac{1}{2} \phi_{\vec{k}}(-\vec{k}) (k^2 + \tau) \phi_{\vec{k}}(\vec{k}) \left[ \begin{array}{l} \vec{k}' = b\vec{k} \\ d^d \vec{k}' = b^d d^d \vec{k} \end{array} \right]$$

$$= \int_0^1 \frac{d^d \vec{k}'}{(2\pi)^d} b^{-d} \frac{1}{2} \phi_{\vec{k}'}(-\vec{k}'/b) (b^{-2} k'^2 + \tau) \phi_{\vec{k}'}(\vec{k}'/b)$$

3. Field renormalization:

with  $\phi'(\vec{k}') = b^y \phi_{\vec{k}'}(\vec{k}'/b)$ :

$$S_{\text{eff}} = \int_0^1 \frac{d^d \vec{k}'}{(2\pi)^d} \frac{1}{2} \phi'(-\vec{k}') (b^{-d-2-2y} k'^2 + b^{-d-2y} \tau) \phi'(\vec{k}')$$

has the same form as original S for  $y = -\frac{d+2}{2}$ .


Then:

$$Z = \int \mathcal{D}\phi' e^{-\int_0^1 \frac{1}{2} \phi'(k'^2 + \tau') \phi'} \quad \text{with } \underline{\underline{\tau' = b^2 \tau}}$$

Beta function ( $0 < \ln b \ll 1$ ):

$$\beta_{\tau} = \frac{d\tau}{d \ln b} = 2\tau \quad (\text{for } u=0)$$

RG flow diagram:



RG for  $\phi^4$  model ( $u > 0$  with  $N=1$ ):

1. Mode elimination:

$$Z = \int \mathcal{D}\phi_{<} \int \mathcal{D}\phi_{>} e^{-S_{0<} - S_{0>} - S_{int}[\phi_{<}, \phi_{>}]}$$

$$= \int \mathcal{D}\phi_{<} e^{-S_{0<}} \int \mathcal{D}\phi_{>} e^{-S_{0>}} \left( 1 - S_{int}[\phi_{<}, \phi_{>}] + \mathcal{O}(u^2) \right)$$



with

$$S_{int}[\phi_{<}, \phi_{>}] = \frac{u}{4!} \left[ \int_0^{\Lambda/b} \int_{\vec{k}_1, \vec{k}_2, \vec{k}_3} \phi_{<} \phi_{<} \phi_{<} \phi_{<} + \int_{\Lambda/b}^{\Lambda} \int_{\vec{k}_1, \vec{k}_2, \vec{k}_3} \phi_{>} \phi_{>} \phi_{>} \phi_{>} + \binom{4}{2} \int_0^{\Lambda} \int_{\vec{k}_1, \vec{k}_2, \vec{k}_3} \phi_{<} \phi_{<} \phi_{>} \phi_{>} \right]$$

$\left[ \int_0^{\Lambda} \int_{\vec{k}} \equiv \int_0^{\Lambda} \frac{d^d k}{(2\pi)^d} \right]$

Thus:

$$Z = Z_{0>} \int \mathcal{D}\phi_{<} e^{-S_{0<}} \left( 1 - \frac{u}{4!} \int_0^{\Lambda/b} \langle \phi_{<} \phi_{<} \phi_{<} \phi_{<} \rangle_{0>} + \int_{\Lambda/b}^{\Lambda} \langle \phi_{>} \phi_{>} \phi_{>} \phi_{>} \rangle_{0>} + \binom{4}{2} \int_0^{\Lambda} \frac{u}{u} \langle \phi_{<} \phi_{<} \phi_{>} \phi_{>} \rangle_{0>} \right) + \mathcal{O}(u^2)$$

 "tree-level"  
 "vacuum"  
 average w.r.t.  $S_{0>}$   
 $\langle \dots \rangle_{0>} \equiv \int \mathcal{D}\phi_{>} e^{-S_{0>}} (\dots) / Z_{0>}$

Wick's theorem:

$$\langle \phi_{<} \phi_{<} \phi_{>} \phi_{>} \rangle_{0>} = \langle \phi_{<} \phi_{<} \rangle_{0>} \langle \phi_{>} \phi_{>} \rangle_{0>} \quad [\text{exercise sheet 3, problem 1c}]$$


Feynman rules (momentum-shell RG):

- vertex  $\times \hat{=} \frac{u}{4!} \delta(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4) (2\pi)^d$
- internal line  $\longleftrightarrow \hat{=} \langle \phi, \phi \rangle_0$
- external line  $\longrightarrow \hat{=} \phi_{\leftarrow}$

Remark:

Only one-particle irreducible (1PI) connected diagrams (that remain connected after cutting one internal line) contribute to the RG flow.

Example:



Reexponentiation:

$$Z = Z_0 \int \mathcal{D}\phi_{\leftarrow} e^{\underbrace{-S_{0\leftarrow} - \frac{u}{4!} \int_0^{\Lambda/b} \phi_{\leftarrow} \phi_{\leftarrow} \phi_{\leftarrow} \phi_{\leftarrow} + \binom{4}{2} \left( \int_0^{\Lambda/b} \phi_{\leftarrow} \phi_{\leftarrow} \right) \left( \int_{\Lambda/b}^{\Lambda} \frac{1}{k^2 + r} \right)}_{=-S_{\text{eff}}}} + \mathcal{O}(u^2)$$

with  $\int_{\Lambda/b}^{\Lambda} \frac{1}{k^2 + r} = \frac{S_d}{(2\pi)^d} \frac{\Lambda^d}{\Lambda^2 + r} \ln b + \mathcal{O}(\ln^2 b)$

2. Momentum rescaling:  $\vec{k}' = b\vec{k}$

$$S_{\text{eff}} = S_{0\leftarrow} + \frac{u}{4!} \left[ \int_0^{\Lambda} b^{-3d} \phi_{\leftarrow} \phi_{\leftarrow} \phi_{\leftarrow} \phi_{\leftarrow} + \binom{4}{2} \int_0^{\Lambda} b^{-d} \phi_{\leftarrow} \phi_{\leftarrow} \frac{S_d}{(2\pi)^d} \frac{\Lambda^d}{\Lambda^2 + r} \ln b \right] + \mathcal{O}(u^2, \ln^2 b)$$

13.5.24

3. Field renormalization:  $\phi'(\vec{k}') = b^{-\frac{d+2}{2}} \phi_{<}(\vec{k}'/b)$

$$S_{eff} = \int_0^\Lambda \frac{1}{2} \phi'(\vec{k}'^2 + \underbrace{b^2 \tau + \frac{u}{2} \frac{S_d}{(2\pi)^d} \frac{\Lambda^d}{\Lambda^{2+\tau}} \ln b}_{\equiv \tau'}) \phi'$$

$$+ \int_0^\Lambda \underbrace{\frac{u}{4!} b^{4-d}}_{\frac{u'}{4!}} \phi' \phi' \phi' \phi' + \mathcal{O}(u^2, \ln^2 b) \quad [b^2 = 1 + \mathcal{O}(\ln b)]$$

Then:

$$\tau' = b^2 \tau + \frac{u}{2} \frac{S_d}{(2\pi)^d} \frac{\Lambda^d}{\Lambda^{2+\tau}} \ln b + \mathcal{O}(u^2, \ln^2 b)$$

$$u' = b^{4-d} u + \mathcal{O}(u^2, \ln^2 b)$$

Introduce dimensionless variables (for convenience):

$$\tau \mapsto t \equiv \frac{\tau}{\Lambda^2}$$

$$u \mapsto g \equiv \frac{S_d}{(2\pi)^d} \frac{u}{\Lambda^{4-d}}$$

Beta functions:

$$\beta_t = \frac{dt}{d \ln b} = 2t + \frac{g}{2} \frac{1}{1+t} + \mathcal{O}(g^2)$$

$$\beta_g = \frac{dg}{d \ln b} = (4-d)g + \mathcal{O}(g^2)$$

Remarks:

• Scaling dimensions  $\dim[\tau] = 2$  and  $\dim[u] = 4-d$  agree with

power-counting dimensions:

$$0 = [S] = \underbrace{[\tau^2]}_{\substack{\text{inverse-length} \\ \text{dimensional} \\ 2}} + [\phi^2] + \underbrace{[d^d x]}_{-d} \Rightarrow [\phi] = \frac{d-2}{2}, \quad [\tau] = 2 = \dim[\tau] \checkmark$$

$$0 = [S] = [u] + \underbrace{[\phi^4]}_{2(d-2)} + \underbrace{[d^d x]}_{-d} \Rightarrow [u] = 4-d = \dim[g] \checkmark$$

• To compute the leading interaction correction to  $\beta_g$  we need to compute the  $g^2$  contribution.



Leading interaction correction to  $\beta_g$  (diagrammatically):



$$= (-1) \frac{1}{2!} \left( \frac{-u}{4!} \right)^2 (\underbrace{\phi\phi\phi\phi}_{\text{re-representation}})(\underbrace{\phi\phi\phi\phi}_{\text{expansion of exp(-)}}) \times \left(\frac{4}{2}\right)^2 \cdot 2$$

where  $\underbrace{\phi\phi}_{\square} \equiv \langle \phi, \phi \rangle_0$

$$= -\frac{1}{4!} \frac{3}{2} u^2 \phi\phi\phi\phi \int_{\Lambda/b}^{\Lambda} \frac{1}{(k^2 + \mu)^2}$$

Thus:

$$\beta_g = (4-d)g - \frac{3}{2} g^2 \frac{1}{(1+t)^2} + \mathcal{O}(g^3)$$

Generalization to  $O(N)$  model:

$$\beta_t = \frac{dt}{d\ln b} = 2t + \frac{N+2}{6} \frac{g}{1+t} + \mathcal{O}(g^2)$$

$$\beta_g = \frac{dg}{d\ln b} = (4-d)g - \frac{N+8}{6} \frac{g^2}{(1+t)^2} + \mathcal{O}(g^3)$$

Fixed points

(a) Gaussian fixed point:  $t^* = g^* = 0$

Near this fixed point:

$\dim[u] = 4-d$  irrelevant for  $d > 4$

$\dim[u_6] = 6-2d$  irrelevant for  $d > 3$

$\uparrow$   
 $\phi^6$  coupling

(b) Wilson-Fisher fixed point: Assume  $\epsilon = 4-d \ll 1$

"fractional dimension"




$$g^* = \frac{6}{N+8} \epsilon + \mathcal{O}(\epsilon^2)$$

$$t^* = -\frac{N+2}{2(N+8)} \epsilon + \mathcal{O}(\epsilon^2)$$

Remark:

Systematic loop expansion: contributions at  $\mathcal{O}(\epsilon^n)$  arise from n-loop Feynman diagrams

# RG flow diagrams:



## Remarks:

- The Gaussian (Wilson-Fisher) fixed governs the critical behavior for  $d > 4$  ( $d < 4$ ).
- $d = d_c^+ = 4$  is the upper critical dimension.

- For  $d > d_c^+$  Landau theory becomes (asymptotically) exact because the theory is effectively Gaussian at criticality.
- An experimental system at  $T_c$  flows to the respective critical fixed point and the system becomes scale invariant.
- The critical behavior is governed by the flow in the vicinity of the critical fixed point.

17.5.24

### Perturbations to the Wilson-Fisher fixed point

Consider RG flow near the WF fixed point:

$$t = t^* + \delta t$$

$$g = g^* + \delta g$$

with  $\delta t \ll t^*$  and  $\delta g \ll g^*$ .

Linearized flow equations ( $O(N)$  model):

$$\frac{d}{d \ln b} \begin{pmatrix} \delta t \\ \delta g \end{pmatrix} = \underbrace{\begin{pmatrix} 2 - \epsilon \frac{N+2}{N+8} & \frac{N+2}{6} \left( 1 + \epsilon \frac{N+2}{2(N+8)} \right) \\ 0 & -\epsilon \end{pmatrix}}_{=: (B_{ij}) \text{ "stability matrix" }} \begin{pmatrix} \delta t \\ \delta g \end{pmatrix} + O(\delta^2)$$

Diagonalization of stability matrix:

$$\sum_{j=1}^2 B_{ij} v_j^I = \theta^I v_i^I \quad I=1,2 \text{ (no sum!)}$$

$\uparrow$  eigenvectors       $\uparrow$  eigenvalues

Remarks:

- $\Theta^I = \text{dim}[v^I]$  is the scaling dimension of the coupling  $v^I$  at the WF fixed point
- Any critical fixed point has exactly one  $\Theta^I > 0$  (w.l.o.g. for  $I=1$ )

Integration of the relevant direction:

$$v^1(b) = v^1(0) b^{\Theta^1} \quad \text{with } \Theta^1 > 0$$

Scaling transformation of reduced temperature  $t_{\text{red}}$  (another tuning parameter):

$$t_{\text{red}} \sim \delta t \sim v^2 \Rightarrow t_{\text{red}} \mapsto b^{\Theta_1} t_{\text{red}} \Rightarrow \Theta_1 \equiv y_t$$

Correlation-length exponent:

$$\boxed{\nu = \frac{1}{\Theta^1}}$$

Wilson-Fisher fixed point:

$$\nu = \frac{1}{2} + \frac{N+2}{4(N+8)} \epsilon + \mathcal{O}(\epsilon^2)$$

Gaussian fixed point:

$$\nu = \frac{1}{2}$$

Remarks:

- For  $N=1$  and  $\epsilon=1$  we get

$$v = \frac{1}{2} + \frac{1}{12} \pm \dots \approx 0.58$$

- Higher-order calculations ( $D=3$ ):


- $v = 0.629(3)$  (six-loop  $\epsilon$  expansion + Borel summation)
- $v = 0.631(4)$  (high-temperature expansion)
- $v = 0.6300(1)$  (MC simulation)
- $v = 0.64(1)$  (neutron scattering of  $FeF_2$ )

[Guida & Zinn-Justin, J.Phys.A 31, 8103 (1998)]

### 4.4 Field-theoretical perspective and anomalous dimension

Idea: Perturbation theory in "renormalized" coupling  $u_R$ :

$$u_R = u - \frac{N+8}{6} u^2 \int_0^\Lambda \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2 + \tau)^2} + O(u^3)$$



Remarks:

- $u_R$  is the effective coupling after integrating out all modes:

- Dimensionless coupling:

$$u \mapsto g \equiv \frac{S_d}{(2\pi)^d} \frac{u}{|\tau|^{(4-d)/2}} \quad \text{diverges for } \tau \rightarrow 0 \text{ when } d < d_c^+ = 4$$

$\Rightarrow$  standard perturbation theory (in  $u$ ) breaks down at criticality

- "Renormalized" perturbation theory (in  $u_R$ ) can be set up to yield finite result.

Example (anomalous dimension, sketch):

Expected critical correlator:


$$\langle \phi(-k) \phi(k) \rangle \propto \frac{1}{k^{2-\eta}}$$

anomalous dimension

Standard perturbation theory:

$$\langle \phi(-k) \phi(k) \rangle \propto \frac{1}{k^2 + \tau - \Sigma(k)}$$

with the "self-energy"



Critical point:  $\tau_R = \tau - \Sigma(0) = 0$

Sunset diagram yields in  $D=4-\epsilon$ :

$$\Sigma(k) - \Sigma(0) = u^2 \left[ c_1 k^2 \ln\left(\frac{\Lambda}{k}\right) + \mathcal{O}(k^4, \epsilon) \right] + \dots$$

constant

To the leading order  $u_R = u + \mathcal{O}(u^2)$  and thus

$$\langle \phi(-k) \phi(k) \rangle \propto \frac{1}{k^2 \left[ 1 + c_2 g_R^2 \ln\left(\frac{\Lambda}{k}\right) \right]} + \mathcal{O}(g_R^3)$$

$$= \frac{1}{k^2} \left( \frac{\Lambda}{k} \right)^{-c_2 g_R^2} + \mathcal{O}(g_R^3)$$

$$\left[ k^x = 1 + x \ln k + \mathcal{O}(x^2) \right]$$

with  $g_R = g^*$  at the critical point.

Reinstating the constants, we read off

(47)

$$\eta = c_2 (g^*)^2 = \frac{N+2}{2(N+8)^2} \varepsilon^2 + \mathcal{O}(\varepsilon^3)$$

Remarks:

- The last step  $1 + c_2 g_R^2 \ln(\frac{\Lambda}{k}) = (\frac{\Lambda}{k})^{c_2 g_R^2} + \mathcal{O}(g_R^4)$  effectively resums an infinite number of diagrams
- For  $N=1$  and  $\varepsilon=1$  (3D Ising):

$$\eta = \frac{1}{54} + \dots \approx 0.02$$

to be compared with (almost exact) value from MC

$$\eta_{MC} = 0.0363(1)$$

31.5.24

## 4.5 Phase transitions and critical dimensions

Universality: different microscopic models flow to the same RG fixed point at criticality

Critical dimensions:

- Upper critical dimension  $d_c^+$ : Mean-field theory asymptotically exact for  $d \geq d_c^+$
- Lower critical dimension  $d_c^-$ : Fluctuations destroy ordered phase at any temperature for  $d \leq d_c^-$
- Critical exponents typically depend on  $d$  for  $d_c^- < d < d_c^+$  and become  $d$ -independent for  $d > d_c^+$  [exception: system with sufficiently long-ranged interactions].

- Classical magnets with short-range interactions [ $O(N)$  models]: (48)

$$d_c^+ = 4 \quad \text{and} \quad d_c^- = \begin{cases} 2 & \text{for } N > 2 \\ 1 & \text{for } N = 1 \end{cases}$$

(The case  $N=2$  and  $d=2$  is special.)

Physics near upper critical dimension [ $O(N)$  models]:

- For  $d < d_c^+ = 4$ : critical fixed point = Wilson-Fisher fixed point,  
observables computable in renormalized perturbation theory in  $u^* = u^*(d)$ ,  
hyperscaling valid.

- For  $d > d_c^+ = 4$ : critical fixed point = Gaussian fixed point,  
observables computable in standard perturbation theory in  $u$ ,  
exponents take mean-field values, e.g.  $\alpha=0$ ,  $\eta=0$ ,  $\nu=\frac{1}{2}$ , etc.  
hyperscaling violated: e.g.

$$2 - \alpha \neq d\nu \quad (\text{Josephson})$$


can be traced back to presence of dangerously irrelevant coupling  $u$   
(free energy nonanalytic at  $u = u^* = 0$ .)

- For  $d = d_c^+ = 4$ : logarithmic corrections to mean-field behavior



# Analytical alternatives to $\epsilon = 4-D$ expansion:

- $\frac{1}{N}$  expansion (exercise sheet 3)
- $2+\epsilon$  expansion: expansion in  $T_c(\epsilon) \propto O(\epsilon)$
- Conformal bootstrap: use symmetry and unitarity arguments to constrain scaling dimensions of operators assuming conformal invariance



world record in precision, e.g.:

$\nu = 0.629971(4)$

(3D Ising, Kos et al., 2016)

[tinyurl.com/ising-bs]

## Summary:

