

Exercises for “Quantum Phase Transitions”

Summer 24

DR. L. JANSSEN

Exercise 1 (19.04.24)

1. Landau functional for a first-order phase transition

(5 pts.)

Consider the free-energy density

$$f(\varphi) = \frac{a}{2}\varphi^2 + \frac{b}{4}\varphi^4 + \frac{c}{6}\varphi^6, \quad (1)$$

which depends on the real order parameter φ . The parameter $a = a(T)$ depends on the temperature T , the coefficients b and c are temperature-independent constants, and $b < 0$, $c > 0$.

- Determine the extrema of the functional $f(\varphi)$ in Eq. (1). List all possibilities and sketch $f(\varphi)$ in each case.
- Calculate the critical value a_c of the parameter a where the position $\varphi_{\text{eq}}(a)$ of the global minimum of $f(\varphi)$ changes discontinuously.
- Sketch the free energy $f(\varphi_{\text{eq}})$ as a function of the parameter a in the vicinity of the phase transition. Why is it a first-order phase transition?

Hint: Expand $f(\varphi_{\text{eq}})$ up to first order in $\delta a = a - a_c$ around $\delta a = 0$.

2. Phase diagram of a two-order-parameter system

(5 pts.)

Consider a system with two real order parameters φ_1 and φ_2 , whose free-energy density is given by

$$f(\varphi_1, \varphi_2) = \frac{r}{2}(\varphi_1^2 + \varphi_2^2) - \frac{g}{2}(\varphi_1^2 - \varphi_2^2) + \frac{u}{4}(\varphi_1^4 + \varphi_2^4) + \frac{v}{2}\varphi_1^2\varphi_2^2, \quad (2)$$

where $u, v > 0$.

- Determine all extrema of the functional $f(\varphi_1, \varphi_2)$ in Eq. (2). Which values are taken by φ_1^2 , φ_2^2 at these extrema?
- Which conditions have to be posed on φ_1^2 and φ_2^2 ? Discuss which phases (i.e., configurations of φ_1 and φ_2) are physically reasonable in which areas of the (r, g) plane.
- In each case, determine the state with the lowest free energy as function of r and g . Distinguish between $u^2 < v^2$ and $u^2 > v^2$.
- Sketch the phase diagram in the (r, g) plane for $u^2 < v^2$ and $u^2 > v^2$, respectively. What are the orders of the different phase transitions?